

DOPbox

Demonstration of the Chebyshev Polynomials

Matthew Harker and Paul O’Leary
Institute for Automation
University of Leoben
A-8700 Leoben, Austria
URL: automation.unileoben.ac.at

Original: January 20, 2013
© 2013

Last Modified: April 11, 2013

Contents

1	Setting things up	2
2	Generate the nodes and basis functions	2
3	Determine the quality of the basis functions	2
4	Plot the basis functions	3
5	Plot the matrix of errors	3

1 Setting things up

Clear up the workspace

```
1 clear;
2 close all;
3 setUpGraphics;
4 %
5 % Define the number of points and the number of basis functions
6 %
7 noPts = 100;
8 noBfs = 5;
```

2 Generate the nodes and basis functions

Generate the Chebyshev nodes

```
9 x = dopNodes( noPts, 'cheby' );
10 %
11 % Setup the first two basis functions
12 %
13 t0 = ones( noPts, 1 );
14 t1 = x;
15 %
16 % Use the Chebyshev recurrence relationship
17 %
18 T = zeros( noPts, noBfs );
19 T(:,1) = t0;
20 T(:,2) = t1;
21 for k=3:noBfs
22     T(:,k) = 2 * x .* T(:,k-1) - T(:,k-2);
23 end;
```

3 Determine the quality of the basis functions

The discrete basis functions should form an orthogonal (but not orthonormal) set of basis functions. The Gram matrix is defined as $\mathbf{G} \triangleq \mathbf{T}^T \mathbf{T}$.

```
24 G = T' * T;
25 %
26 % Removing the diagonal elements should yield a matrix of
27 % zeros.
28 %
29 Gperp = diag(diag(G)) - G;
30 %
31 % The Frobenius norm of Gperp is now used as a measure of quality for
    the
```

```

32 % basis functions.
33 %
34 e = (norm( Gperp, 'fro' ));
    and the  $n_d = -\log_{10}$  as an estimate for the number of available digits.
35 nd = - log10( e );

```

4 Plot the basis functions

Plot the Chebyshev basis functions

```

36 figure
37 plot( x, t1, 'k' );
38 hold on;
39 for k=2:noBfs
40     plot( x, T(:,k), 'k' );
41 end;
42 grid on;
43 xlabel('Support');
44 ylabel('Value');
45 title(['The first n = ',int2str( noBfs ),' Chebyshev basis functions (
        no digits n_d = ',num2str(nd),')']);

```




Figure 1: Some Chebyshev polynomials.

5 Plot the matrix of errors

Plot the error matrix.

```

46 fig2 = figure;
47 imagesc( Gperp );
48 colorbar;
49 axis image;
50 xlabel('Basis function no. ');
51 ylabel('Basis function no. ');

```




Figure 2: Projection of the basis functions onto their orthogonal complement.